



# basic education

Department:  
Basic Education  
**REPUBLIC OF SOUTH AFRICA**

## NATIONAL SENIOR CERTIFICATE NASIONALE SENIOR SERTIFIKAAT

**GRADE/GRAAD 12**

**PHYSICAL SCIENCES: CHEMISTRY (P2)  
FISIESE WETENSKAPPE: CHEMIE (V2)**

**NOVEMBER 2019**

**MARKING GUIDELINES/NASIENRIGLYNE**

**MARKS/PUNTE: 150**

**DEPARTMENT OF BASIC  
EDUCATION**

PRIVATE BAG X895, PRETORIA 0001

2019 -11- 13

**APPROVED MARKING GUIDELINE**

**PUBLIC EXAMINATION**

**These marking guidelines consist of 20 pages.  
Hierdie nasienriglyne bestaan uit 20 bladsye.**

Approved  
M Edelos  
DBE CE  
13/11/2019

Approved  
Flower  
DBE FM  
13/11/2019

approved  
DBE Pnt. Mod.  
13/11/2019

-Approved  
Sibani  
13/11/2019

Approved!  
Vinh  
13/11/2019

**QUESTION 1/VRAAG 1**

- 1.1 D ✓✓ (2)
  - 1.2 C ✓✓ (2)
  - 1.3 B ✓✓ (2)
  - 1.4 D ✓✓ (2)
  - 1.5 C ✓✓ (2)
  - 1.6 B ✓✓ (2)
  - 1.7 B ✓✓ (2)
  - 1.8 A ✓✓ (2)
  - 1.9 A ✓✓ (2)
  - 1.10 C ✓✓ (2)
- [20]**


**QUESTION 2/VRAAG 2**

2.1

2.1.1  $C_nH_{2n-2}$  ✓


2.1.2

*refer for dummy ①*



**Marking criteria/Nasiennriglyne**

- Functional group correct. ✓  
*Funksionele groep korrek.*
- 2 methyl substituents. ✓  
*2 metielsubstituente.*
- Whole structure correct. *Hele struktuur korrek:* 3/3



DEPARTMENT OF BASIC EDUCATION  
 PRIVATE BAG X895, PRETORIA 0001  
 2019 -11- 13  
 APPROVED MARKING GUIDELINE  
 PUBLIC EXAMINATION

*Handwritten signatures and initials.*

2.2

2.2.1 Compounds with the same molecular formula, ✓ but different positions of the side chain/substituents/functional groups ✓ on the parent chain.  
*Verbindings met dieselfde molekulêre formule, maar verskillende posisies van die syketting/substituente/funksionele groepe op die stamketting.* (2)

2.2.2 Pentan-3-one/3-pentanone ✓✓  
 Pentan-3-oon/3-pentanoon

butan-3-one ✓

**Marking criteria/Nasienriglyne**

- Functional group and correct position i.e. 3 /Funksionele groep en korrekte posisie nl. 3. ✓
- Whole name correct/Hele naam korrek. ✓


**Accept for ONE mark/Aanvaar vir EEN punt** ✓ 1/2

Pentanone with the 3 in incorrect place, e.g. penta-3-none.  
 Pentanoon met die 3 in foutiewe plek, bv. penta-3-noon.

pent-3-one ✓

(2)

2.2.3




**Marking criteria/Nasienriglyne**


- Whole structure correct:/Hele struktuur korrek: 2/2
- Only functional group correct/Slegs funksionele groep korrek Max: 1/2

**OR: Any correct structure of an aldehyde with five carbon atoms.**

**OF: Enige korrekte struktuur van 'n aldehied met vyf koolstofatome.**



(2)



*Handwritten signatures and initials.*

2.3  
2.3.1 Tertiary (alcohol)/Tersiêre (alkohol) ✓

The C atom bonded to the functional group/hydroxyl (group)/-OH is bonded to three other C atoms. /The C-atom bonded to the hydroxyl (group) has no hydrogen atoms. ✓

Die C-atoom gebind aan die funksionele groep/hidroksiel(groep)/-OH is gebind aan drie ander C-atome./ Die C-atoom gebind aan die hidroksiel (groep) het geen waterstofatome nie.

(2)

2.3.2 2-methylbutan-2-ol/2-methyl-2-butanol/2-metielbutan-2-ol/2-metiel-2-butanol

**Marking criteria/Nasienriglyne**

- 2-methyl/2-metiel ✓
- Butan-2-ol/2-butanol ✓
- Any error e.g. hyphens omitted and/or incorrect sequence:

Enige fout, bv. koppeltekens weggelaat en/of verkeerde volgorde: Max./Maks:  $\frac{1}{2}$

(2)

2.3.3 2-methylbut-2-ene/2-methyl-2-butene/2-metielbut-2-een/2-metiel-2-buteen

**Marking criteria/Nasienriglyne**

- 2-methyl/2-metiel ✓
- But-2-ene/2-butene/But-2-een/2-buteen ✓
- Any error e.g. hyphens omitted and/or incorrect sequence:

Enige fout, bv. koppeltekens weggelaat en/of verkeerde volgorde: Max./Maks:  $\frac{1}{2}$

(2)

[16]

QUESTION 3/VRAAG 3

3.1 **Marking guidelines/Nasienriglyne**

The underlined key phrases must be used in the **CORRECT CONTEXT (pressure/boiling)**. /Die onderstreepte frases moet gebruik word in die **KORREKTE KONTEKS (druk/kook)**.

The temperature ✓ at which the vapour pressure of a substance equals atmospheric/external pressure. ✓

Die temperatuur waar die dampdruk van 'n stof gelyk is aan atmosferiese/eksterne druk.

(2)


3.2 (Q, R and S) have same molecular mass/formulae/number of carbon and hydrogen atoms/are (chain) isomers. ✓ same no of C atoms and same homologous series  
(Q, R en S) het dieselfde molekulêre massa/formule/aantal koolstof en waterstofatome/ is (ketting)isomere.

OR/OF

The compounds are all alkanes /same homologous series and have the same number of carbon atoms.

Die verbinding is almal alkane /dieselfde homoloë reeks en het die dieselfde aantal koolstofatome.

(1)



**Marking guidelines/Nasienriglyne**

- 55 (°C) ✓
- Compare all three compounds or Q and S in terms of branches/chain lengths / surface area. ✓  
*Vergelyk al drie verbindings of Q en S in terme van vertakkings/kettinglengte/ oppervlakarea.*
- Compare strengths of all three or Q and S's IMF's / Vergelyk sterkte van al drie of Q en S se IMK'e. ✓
- Compare energy of all three / Vergelyk energie van al drie. ✓

3.3

55 (°C) ✓

**Compare compound R with compounds Q and S:**

- Compound R is less branched/compact/spherical/surface area than compound Q and more branched/compact/spherical/surface area than compound S. ✓  
**OR**  
Q is the most branched/compact /spherical/surface area and S is least branched/compact/spherical/surface area.
- Intermolecular forces in compound R are stronger than in compound Q and weaker than in compound S. ✓
- More energy needed to overcome intermolecular forces in compound R than in compound Q and less energy needed to overcome (break) intermolecular forces in compound R than in compound S. ✓

**OR**

- Compound R has a longer chain length than compound Q and a shorter chain length than compound S. ✓  
**OR**  
S has the longest chain length and Q the shortest.
- Intermolecular forces increase with increase in chain length. ✓
- More energy needed to overcome intermolecular forces as chain length increases. ✓

**Vergelyk verbinding R met verbindings Q en S:**

Verbinding R is minder vertak/kompak/sferies/oppervlak as verbinding Q en meer vertak as verbinding S.

**OF**


Q is die meeste vertak/kompak en S is die minste vertak/kompak/series/oppervlak.

Intermolekulêre kragte in verbinding R is sterker as in verbinding Q en swakker as in verbinding S.

Meer energie word benodig om intermolekulêre kragte in verbinding R te oorkom as in verbinding Q, en minder energie word benodig om intermolekulêre kragte in verbinding R te oorkom / breek as in verbinding S.

**OF**

- Verbinding R het 'n langer kettinglengte as verbinding Q en 'n korter kettinglengte as S.
- **OR**  
S het die langste ketting en Q die kortste.
- Intermolekulêre kragte neem toe met toename in kettinglengte.
- Meer energie word benodig om intermolekulêre kragte te oorkom wanneer kettinglengte toeneem.



independent  
ask for  
Q short and  
S long

R longer than Q  
Q shorter than S

Q long  
Q shorter  
Q shortest

Q longest  
Q shorter than S.  
Q shortest.

Handwritten signature and initials.

3.4

3.4.1 P ✓✓

(2)

3.4.2

**Marking guidelines/Nasienriglyne**

- Name type of IMFs in P/pentanal. ✓  
*Noem tipe IMK'e in P/pentanaal.*
- Name type of IMFs in/Noem tipe IMK'e in T/pentan-1-ol. ✓
- Compare strength of IMFs. /Vergelyk sterkte van IMK'e. ✓  
**OR/OF**  
Compare energy needed to overcome IMFs./Vergelyk energie benodig om IMK'e te oorkom.

- In P/ pentanal/aldehydes: dipole-dipole forces. ✓ (in addition to London forces/dispersion forces/induced dipole forces). *— not H-forces.*
- In T/pentan-1-ol: Hydrogen bonding. ✓ (in addition to London forces/dispersion forces/induced dipole forces).
- Intermolecular forces in P/pentanal are weaker. ✓ than in T/pentan-1-ol **OR** dipole-dipole forces are weaker than hydrogen bonds **OR** intermolecular forces in T/pentan-1-ol are stronger than in P/pentanal. **OR**  
More energy needed to overcome/break intermolecular forces in T.
- In P/pentanaal/aldehyede: dipool-dipoolkragte (tesame met Londonkragte/dispersiekragte/geïnduseerde dipoolkragte).
- In T/pentan-1-ol: Waterstofbinding. (tesame met Londonkragte/dispersiekragte/geïnduseerde dipoolkragte).
- Intermolekulêre kragte in P swakker as in T/pentan-1-ol **OF** intermolekulêre kragte in T/pentan-1-ol sterker as in P/pentanaal **OF** dipool-dipoolkragte is swakker as waterstofbindings. **OF**  
Meer energie benodig om intermolekulêre kragte te oorkom/breek in T.

(3)  
[12]




*Handwritten signatures and initials.*

Alex 4541002  
4341032 (320136)


## QUESTION 4/VRAAG 4

- 4.1 Haloalkane/alkyl halide ✓  
Haloalkaan/alkielhalied (1)
- 4.2 Elimination/dehydrohalogenation ✓  
Eliminasie/dehidrohalogenering (1)
- 4.2.2 Substitution/hydrolysis ✓  
Substitusie/hidrolise (1)
- 4.2.3 Esterification/condensation ✓  
Esterifikasie/kondensasie/verestering (1)
- 4.3  
4.3.1 • (Mild) heat/Heating/(matige) hitte/verhitting ✓  
• Dilute (strong base)/Verdunde (sterk basis)/(NaOH/KOH/LiOH) ✓  
OR/OR  
• Add water/H<sub>2</sub>O/Voeg water/H<sub>2</sub>O by  
strong heat x  
Dilute acid x (2)
- 4.3.2 Propan-1-ol/1-propanol ✓✓

**Marking criteria/Nasienriglyne:**

- Correct stem and functional group i.e. propanol/Korrekte stam en funksionele groep, d.i. propanol. ✓
  - Whole name correct:/Hele naam korrek: propan-1-ol ✓
- prop-1-ol x

4.4

**Marking criteria/Nasienriglyne**

- Whole structure correct:/Hele struktuur korrek: 2/2
- Only functional group correct/Slegs funksionele groep korrek: 1/2

**Notes/Aantekeninge**


- Accept -OH as condensed. /Aanvaar -OH as gekondenseerd.
- Condensed or semi-structural formula:  
Gekondenseerde of semi-struktuurformule: Max./Maks. 1/2
- Molecular formula/Molekulêre formule: 0/2
- If functional group is incorrect/Indien funksionele groep verkeerd is: 0/2
- If more than one functional group:  
Indien meer as een funksionele groep: 0/2

Handwritten signatures and initials at the bottom right of the page.

4.5

**POSITIVE MARKING FROM Q4.3.2 ONLY IF THE COMPOUND IN Q4.3.2 IS AN ALCOHOL. /POSITIEWE NASIEN VANAF V4.3.2 SLEGS INDIEN DIE VERBINDING IN Q4.3.2 'N ALKOHOL IS.**

4.5.1



**Marking criteria/Nasienriglyne**

- Whole structure correct:/Hele struktuur korrek: 2/2
- Only functional group correct/Slegs funksionele groep korrek: 1/2

**Notes/Aantekeninge**

- Condensed or semi-structural formula:  
 Gekondenseerde of semistruktuurformule: Max./Maks. 1/2
- Molecular formula/Molekulêre formule: 0/2
- If functional group is incorrect/Indien funksionele groep verkeerd is: 0/2

4.5.2 (Concentrated) sulphuric acid/(Gekonsentreerde) swawelsuur/H<sub>2</sub>SO<sub>4</sub> ✓ (1)  
 Hydrogen sulfate refer [13]

**QUESTION 5/VRAAG 5**

5.1 Exothermic/Eksotermies ✓


⊖ ΔH < 0/Energy is released/Energie word vrygestel ✓ Energy is one of the products (2)

5.2

2 - 0,25 ✓  
 30 ✓  
 or  
 2 - 0,25 ✓  
 0 - 30 ✓

$\text{rate/tempo} = -\frac{\Delta m}{\Delta t}$ $= -\frac{0,25 - 2}{30} \checkmark$ $= 0,06 \text{ (g} \cdot \text{s}^{-1}) \checkmark$ $\text{(0,0583 g} \cdot \text{s}^{-1})$	<p><b>OR/OF</b></p> $\text{rate/tempo} = -\frac{\Delta m}{\Delta t}$ $\geq -\frac{-1,75}{30} \checkmark$ $= 0,06 \text{ (g} \cdot \text{s}^{-1}) \checkmark$ $\text{(0,0583 g} \cdot \text{s}^{-1})$
--	--

**Notes/Aantekeninge**  
 (Accept) negative answer i.e./Aanvaar negatiewe antwoord d.i. - 0,06 g · s<sup>-1</sup>.



Handwritten signatures and initials.



5.3

<b>Marking guidelines</b>	
<ul style="list-style-type: none"> <li>Calculate/Bereken: <math>m(\text{CaCO}_3)</math> reacted/reageer or <math>l</math> of <math>V(\text{CO}_2)</math> produced/gevorm. ✓</li> <li>Substitute/Vervang: <math>100 \text{ g} \cdot \text{mol}^{-1}</math>. ✓</li> <li>USE mol ratio/GEBRUIK molverhouding: <math>n(\text{CO}_2) : n(\text{CaCO}_3) = 1 : 1</math> ✓</li> <li>Use of / Gebruik van <math>22,4 \text{ dm}^3 \cdot \text{mol}^{-1}</math>. ✓</li> <li>Final answer/Finale antwoord: <math>0,18 \text{ dm}^3</math> (<math>0,1792 \text{ dm}^3</math>) ✓</li> </ul>	
<b>OPTION 1/OPSIE 1</b>	<b>OPTION 2/OPSIE 2</b>
$m(\text{CaCO}_3) = \frac{40}{100} \times 2 \checkmark$ $= 0,8 \text{ g}$ $n(\text{CaCO}_3)_{\text{reacted}} = \frac{m}{M}$ $= \frac{0,8}{100} \checkmark$ $= 8 \times 10^{-3} \text{ mol}$ <p><i>(Handwritten: <math>40+12+3 \cdot 16</math>)</i></p> $n(\text{CO}_2) = n(\text{CaCO}_3) \checkmark$ $= 8 \times 10^{-3} \text{ mol}$ $V(\text{CO}_2) = 8 \times 10^{-3} \times 22,4 \checkmark$ $= 0,18 \text{ dm}^3 \checkmark$	<b>For 2 g antacid/teensuurtablet:</b> $100 \text{ g} \checkmark \text{CaCO}_3 \dots\dots 22,4 \text{ dm}^3 \checkmark \text{CO}_2$ $2 \text{ g CaCO}_3 \dots\dots 0,448 \text{ dm}^3 \checkmark$  $100\% \text{CO}_2 \dots\dots 0,448 \text{ dm}^3 \checkmark$ $40\% \text{CO}_2 \dots\dots 0,18 \text{ dm}^3 \checkmark$
	<b>OPTION 3/OPSIE 3</b>
	$100\% \text{CaCO}_3 \dots\dots 2 \text{ g}$ $40\% \dots\dots 0,8 \text{ g} \checkmark$  $100 \text{ g} \checkmark \dots\dots 1 \text{ mol}$ $0,8 \text{ g} \dots\dots 8 \times 10^{-3} \text{ mol} \checkmark$  $1 \text{ mol} \dots\dots 22,4 \text{ dm}^3 \checkmark$ $8 \times 10^{-3} \text{ mol} \dots\dots 0,18 \text{ dm}^3 \checkmark$

(5)

5.4

**ANY ONE/ENIGE EEN:**


- Concentration (of acid)/Konsentrasie (van suur) ✓
- Size/mass of tablet/Identical tablet /Type of tablet. *or mass of  $\text{CaCO}_3$  / mass of antacid ✓*  
 Grootte/massa van tablet/Identiese tablet./Tipe tablet.
- State of division / Surface area / Toestand van verdeeldheid / reaksieoppervlak.

(1)

5.5

*see variables correct relation ✓*

<b>Criteria for conclusion/Riglyne vir gevolgtrekking:</b>	
Dependent [(reaction) rate/time] and independent (temperature) variables correctly identified.	✓
Afhanklike [(reaksie)tempo/tyd] en onafhanklike (temperatuur) veranderlikes korrek geïdentifiseer.	✓
Relationship between the independent and dependent variables correctly stated./Verwantskap tussen die afhanklike en onafhanklike veranderlikes korrek genoem.	✓



*Handwritten signatures and initials*

**Examples/Voorbeelde:**

- Reaction rate ( $\frac{1}{\text{time}}$ ) increases with increase in temperature.  
*Reaksietyempo ( $\frac{1}{\text{time}}$ ) neem toe met toename in temperatuur.*
- Reaction rate ( $\frac{1}{\text{time}}$ ) decreases with decrease in temperature.  
*Reaksietyempo ( $\frac{1}{\text{time}}$ ) neem af met afname in temperatuur.*
- Time taken for reaction decreases when temperature increases.  
*Tyd vir die reaksie neem af wanneer temperatuur toeneem.*
- Time taken for reaction increases when temperature decreases.  
*Tyd vir die reaksie neem toe as temperatuur afneem.*

**IF/INDIEN**

Reaction rate is DIRECTLY proportional to temperature: Max.  $\frac{1}{2}$   
*Reaksietyempo is DIREK eweredig aan temperatuur: Maks.  $\frac{1}{2}$*

(2)

5.6

*speed higher ✓*


- Increase in temperature increases the average kinetic energy/molecules move faster. / *Toename in temperatuur verhoog die gemiddelde kinetiese energie/molekule beweeg vinniger. ✓*
- More molecules have enough/sufficient kinetic energy/More molecules have  $E_k > E_a$ . ✓  
*Meer molekule het genoeg/voldoende kinetiese energie/Meer molekule het  $E_k > E_a$ .*
- More effective collisions per unit time/second. / Frequency of effective collisions increases. ✓  
*Meer effektiewe botsings per eenheidtyd/sekonde./Frekwensie van effektiewe botsings neem toe.*

(3)

5.7

**Marking guidelines/Nasienriglyne**

- For each value of temperature, the CURVE Y must be above the given CURVE. / *Vir elke waarde van temperatuur, moet kurwe Y bo die gegewe kurwe wees. ✓*
- CURVE Y must have an increasing rate with an increase in temperature. / *KURWE Y moet 'n toenemende tempo het soos die temperatuur toeneem. ✓*



DEPARTMENT OF BASIC EDUCATION  
 PRIVATE BAG 2855, PRETORIA 0001  
 2019-11-13  
 APPROVED MARKING GUIDELINE  
 PUE LIC EXAMINATION

*Van. ✓*  
*EW*  
*Sub.*

(2)  
 [18]

### QUESTION 6/VRAAG 6

- 6.1 (The stage in a chemical reaction when the) rate of forward reaction equals the rate of reverse reaction. ✓✓  
(Die stadium in 'n chemiese reaksie wanneer die) tempo van die voorwaartse reaksie gelyk is aan die tempo van die terugwaartse reaksie. (2 or/of 0)

#### OR/OF


(The stage in a chemical reaction when the) concentrations of reactants and products remain constant.

(Die stadium in 'n chemiese reaksie wanneer die) konsentrasies van reaktanse en produkte konstant bly. (2 or/of 0) (2)

### 6.2 CALCULATIONS USING NUMBER OF MOLES BEREKENINGE WAT AANTAL MOL GEBRUIK

#### 6.2.1 Marking guidelines/Nasienriglyne

- Substitute/Vervang: 44 g·mol<sup>-1</sup>. ✓
- Equilibrium concentration of CO<sub>2</sub> multiply by 3 dm<sup>3</sup>  
Ewewigskonsentrasie van CO<sub>2</sub> vermenigvuldig met 3 dm<sup>3</sup> } ✓  
AND/EN n(CO)<sub>eq</sub> divide by /deel deur 3 dm<sup>3</sup>
- Use mole ratio/Gebruik molverhouding: 1:2 / n(CO) = 2n(CO<sub>2</sub>). ✓
- $n(\text{CO}_2)_{\text{change}} = n(\text{CO}_2)_{\text{initial}} - n(\text{CO}_2)_{\text{final}}$  } ✓  
 $n(\text{CO})_{\text{eq/ewe}} = n(\text{CO})_{\text{initial/begin}} + \Delta n(\text{CO})$  }
- Correct K<sub>c</sub> expression (formulae in square brackets). ✓  
Korrekte K<sub>c</sub>-uitdrukking (formules in vierkanthakies).
- Substitution of concentrations into K<sub>c</sub> expression. ✓  
Vervanging van konsentrasies in K<sub>c</sub>-uitdrukking.
- Final answer/Finale antwoord: 12,24 (range/gebied: 11,85 – 12,66) ✓



0720203713  
Sib. Mr. J. M. J. M.

**OPTION 1/OPSIE 1**

$$n(\text{CO}_2) = \frac{m}{M}$$

$$= \frac{60,8}{44} \checkmark (4)$$

$$= 1,382 \text{ mol}$$

	CO <sub>2</sub>	CO
Initial quantity (mol) Aanvangs hoeveelheid (mol)	1,382	0
Change (mol) Verandering (mol)	1,22	2,44
Quantity at equilibrium (mol)/ Hoeveelheid by ewewig (mol)	0,162	2,44
Equilibrium concentration (mol·dm <sup>-3</sup> ) Ewewigskonsentrasie (mol·dm <sup>-3</sup> )	0,054	0,813

Use ratio/Gebruik verhouding  $\checkmark (5)$   
 and  
 Divide/multiply by 3/Deel/vermenigvuldig met 3  $\checkmark (7)$

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \checkmark (1)$$

$$= \frac{(0,813)^2}{0,054} \checkmark (2)$$

$$= 12,24 \checkmark (3)$$

if wrong, then stop marking.

No K<sub>c</sub> expression, correct substitution/Geen K<sub>c</sub>-uitdrukking, korrekte substitusie: Max./Maks.  $\frac{6}{7}$   
 Wrong K<sub>c</sub> expression/Verkeerde K<sub>c</sub>-uitdrukking: Max./Maks.  $\frac{4}{7}$

**OPTION 2/OPSIE 2**

$$n(\text{CO}_2) = \frac{m}{M}$$

$$= \frac{60,8}{44} \checkmark$$

$$= 1,382 \text{ mol}$$

$$n(\text{CO}_2)_{\text{change}} = n(\text{CO}_2)_{\text{initial/begin}} - n(\text{CO}_2)_{\text{final/finaal}}$$

$$= 1,382 - 0,162$$

$$= 1,22 \text{ mol}$$

$$n(\text{CO})_{\text{change}} = 2(\text{CO}_2)_{\text{change}} \checkmark$$

$$= 2(1,22) \checkmark$$

$$= 2,44 \text{ mol}$$

$$n(\text{CO})_{\text{eq}} = n(\text{CO})_{\text{change}} = 2,44 \text{ mol}$$

$$c(\text{CO}) = \frac{n}{V}$$


$$= \frac{2,44}{3} \checkmark$$

$$= 0,813 \text{ mol·dm}^{-3}$$

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \checkmark$$

$$= \frac{(0,813)^2}{0,054} \checkmark$$

$$= 12,24 \checkmark \text{ (Accept range/Aanvaar gebied: 11,85 – 12,66).}$$



Handwritten signatures and initials.

**CALCULATIONS USING CONCENTRATION**  
**BEREKENINGE WAT KONSENTRASIE GEBRUIK**

**Marking guidelines/Nasienriglyne**

- Substitute 44 g·mol<sup>-1</sup>. ✓
- Initial n(CO<sub>2</sub>) divide by 3 dm<sup>3</sup>. ✓  
 Aanvanklike n(CO<sub>2</sub>) gedeel deur 3 dm<sup>3</sup>.
- **USE** ratio/**GEBRUIK** verhouding: c(CO<sub>2</sub>) : c(CO) = 1 : 2 ✓
- $\Delta c(\text{CO}_2) = c(\text{CO}_2)_{\text{initial/begin}} - c(\text{CO}_2)_{\text{eq/ewe}}$ .  
 $c(\text{CO})_{\text{eq/ewe}} = c(\text{CO})_{\text{initial/begin}} + \Delta c(\text{CO})$ . } ✓
- Correct K<sub>c</sub> expression (formulae in square brackets). ✓  
 Korrekte K<sub>c</sub> uitdrukking (formules in vierkanthakies).
- Substitution of concentrations into K<sub>c</sub> expression. ✓  
 Vervanging van konsentrasies in K<sub>c</sub>-uitdrukking.
- Final answer/Finale antwoord: 12,15 (range/gebied: 11,85 – 12,66) ✓

4/7

DEPARTMENT OF BASIC EDUCATION  
 PRIVATE BAG 9373, PRETORIA 0001  
 2019-11-13  
 APPROVED MARKING GUIDELINE  
 PUF LIC EXAMINATOR

**OPTION 3/OPSIE 3**

$$n(\text{CO}_2) = \frac{m}{M}$$

$$= \frac{60,8}{44} \checkmark$$

$$= 1,382 \text{ mol}$$

	CO <sub>2</sub>	CO
Initial concentration (mol·dm <sup>-3</sup> ) Aanvanklike konsentrasie (mol·dm <sup>-3</sup> )	0,4607	0
Change (mol·dm <sup>-3</sup> ) Verandering (mol·dm <sup>-3</sup> )	0,4067	0,813
Equilibrium concentration (mol·dm <sup>-3</sup> ) Ewewigskonsentrasie (mol·dm <sup>-3</sup> )	0,054	0,813

Divide by /Deel deur 3 dm<sup>3</sup> ✓  
 ratio ✓  
 verhouding

$$K_c = \frac{[\text{CO}]^2}{[\text{CO}_2]} \checkmark$$

$$= \frac{(0,813)^2}{0,054} \checkmark$$

$$= 12,15 \checkmark$$

No K<sub>c</sub> expression, correct substitution/Geen K<sub>c</sub>-uitdrukking, korrekte substitusie: Max./Maks. 6/7

Wrong K<sub>c</sub> expression/Verkeerde K<sub>c</sub>-uitdrukking: Max./Maks. 4/7

(7)

**6.2.2 POSITIVE MARKING FROM Q6.2.1/POSITIEWE NASIEN VANAF V6.2.1**

$$n(\text{C})_{\text{reacted/reageer}} = n(\text{CO}_2)_{\text{reacted/reageer}}$$

$$= 1,22 \text{ mol} \checkmark$$

from table

$$m(\text{C}) = nM$$

$$= 1,22(12) \checkmark$$

$$= 14,64 \text{ g} \checkmark$$

**Marking guidelines**

- **USE** mol ratio/ **GEBRUIK** molverhouding: n(C) = n(CO<sub>2</sub>). ✓
- Substitute/Vervang: 12 g·mol<sup>-1</sup>. ✓
- Final answer/Finale antwoord: 14,64 g. ✓

(3)

Not:

$$c = \frac{m}{MV}$$

$$0,406 = \frac{m}{12(3)}$$

$$m = 14,62 \text{ g.}$$

C is solid, does not change conc.  
 Copyright reserved/Kopiereg voorbehou

Handwritten signatures and marks.

6.3

6.3.1 Remains the same/Bly dieselfde ✓

(1)

6.3.2 Decreases/Afneem ✓ (amount of CO will decrease)

- (When pressure is increased) the reaction that leads to the smaller amount/number of moles/volume of gas is favoured. ✓  
(Wanneer die druk verhoog word,) word die reaksie wat tot die kleiner hoeveelheid/aantal mol/volume gas lei, bevoordeel.
- The reverse reaction is favoured. / More CO<sub>2</sub> is formed. ✓  
Die terugwaartse reaksie word bevoordeel./ meer CO<sub>2</sub> word gevorm.

(3)

6.4

6.4.1 Endothermic/Endotermies ✓

- When the temperature increases the mol/percentage CO(g)/product increases/forward reaction is favoured./Wanneer die temperatuur toeneem, neem die mol/persentasie CO(g)/produk toe/voorwaartse reaksie word bevoordeel. ✓
- An increase in temperature favours the endothermic reaction./Toename in temperatuur bevoordeel die endotermiese reaksie. ✓

(3)

6.4.2

**POSITIVE MARKING FROM Q6.2.1./POSITIEWE NASIEN VANAF V6.2.1.**

**Marking guidelines/Nasienriglyne**

- Calculate total volume/mol of gas at equilibrium/Bereken totale volume/mol gas by ewewig:  $0,162 + 2,44 = 2,606 \text{ dm}^3 / \text{mol}$  ✓  
**OR/OF**  
Calculate the total concentration at equilibrium/Bereken die totale konsentrasie by ewewig:  $0,054 + 0,813 = 0,867 \text{ mol} \cdot \text{dm}^{-3}$
- Calculate percentage of ANY one gas/Bereken persentasie van ENIGE een gas (CO<sub>2</sub> or/of CO). ✓
- Final answer/Finale antwoord:  $T = 827 \text{ }^\circ\text{C}$  ✓


Dummy 3: Calculate % for volume / mole / conc NOT mass

**OPTION 1/OPSIE 1**

$n_{\text{total eq}} = 0,162 + 2,44 \checkmark \rightarrow m = nM$   
 $= 2,606 \text{ dm}^3$   
 $\% \text{ CO}_2 = \frac{0,162}{2,606} \times 100 \checkmark$  **OR/OF**  $\% \text{ CO} = \frac{2,44}{2,606} \times 100 \checkmark$   
 $= 6,225 \%$   $= 93,63 \%$

**OPTION 2/OPSIE 2**

$n = cV \rightarrow m = nM$   
 $c_{\text{total eq}} = 0,054 + 0,813$   
 $= 0,867 \text{ mol} \cdot \text{dm}^{-3}$  } using masses 3/3  
 add masses }  
 get % }  
 $\% \text{ CO}_2 = \frac{0,054}{0,867} \times 100 \checkmark$  **OR/OF**  $\% \text{ CO} = \frac{0,813}{0,867} \times 100 \checkmark$   
 $= 6,228 \%$   $= 93,77 \%$



∴  $T = 827 \text{ }^\circ\text{C}$  ✓ Independent.

(3)  
[22]

**QUESTION 7/VRAAG 7**

7.1 Strong (acid)/Sterk (suur) ✓



Large/Groot  $K_a$  value/waarde/  $K_a > 1$  / (HBr) ionises completely/ioniseer volledig ✓

(2)

7.2  $H_2O$  ✓

$Br^-$  ✓

(2)

7.3

7.3.1

**Marking guidelines/Nasienriglyne**

- Formula/Formule:  $c = \frac{n}{V} / n = cV / \frac{c_a \times V_a}{c_b \times V_b} = \frac{n_a}{n_b}$  ✓
- Substitution of/Vervanging van:  $(0,5)(0,0165)/(0,5)(16,5)$  ✓
- Use mol ratio/Gebruik molverhouding:  $1:1/n(HBr) = n(NaOH)$  ✓
- Substitute/Vervang:  $V = 0,09 \text{ dm}^3 / 90 \text{ cm}^3$  ✓
- Formula/Formule:  $pH = -\log[H_3O^+]$  ✓
- Substitute  $[H_3O^+]$  in pH formula. ✓
- Final answer/Finale antwoord:  $pH = 1,04$  (range/gebied:  $1,036 - 1,05$ ) ✓

**OPTION 1/OPSIE 1**

$$n(NaOH)_{\text{reacted/reageer}} = cV \checkmark$$

$$= 0,5(0,0165) \checkmark$$

$$= 0,00825 \text{ mol}$$

$$n(HBr)_{\text{excess/oormaat}} = n(NaOH) = 0,00825 \text{ mol} \checkmark$$

$$c(H_3O^+) = \frac{n}{V}$$

$$= \frac{0,00825}{0,09} \checkmark$$

$$= 0,092 \text{ mol} \cdot \text{dm}^{-3}$$

$$pH = -\log[H_3O^+] \checkmark$$

$$= -\log(0,092) \checkmark$$

$$= 1,04 \checkmark$$

Carry down whatever they have for  $C_a$

**OPTION 2/OPSIE 2**

$$\frac{c_a V_a}{c_b V_b} = \frac{n_a}{n_b} \checkmark$$

$$\frac{c_a (90)}{(0,5)(16,5)} \checkmark = \frac{1}{1} \checkmark$$


$$C_a = 0,092 \text{ mol} \cdot \text{dm}^{-3}$$

$$pH = -\log[H_3O^+] \checkmark$$

$$= -\log(0,092) \checkmark$$

$$= 1,04 \checkmark$$

(7)



Handwritten signatures and initials.

*must be together*

7.3.2

**Marking guidelines/Nasienriglyne**

- Calculate/Bereken  $n(\text{HBr})_{\text{initial/aanvanklik}}$ : substitute/vervang  $(0,45)(0,09)$  in  $n = cV$  ✓
- Subtraction/Aftrekking:  
 $n(\text{HBr})_{\text{reacted/reageer}} = n(\text{HBr})_{\text{initial/aanvanklik}} - n(\text{HBr})_{\text{reacted with/reageer met NaOH}}$  ✓✓  
*1 mark for subtract, then rest for all*
- OR/OF:  $c(\text{HBr})_{\text{reacted/reageer}} = c(\text{HBr})_{\text{initial/aanvanklik}} - c(\text{H}_3\text{O}^+)_{\text{excess/oormaat}}$  ✓
- Use mol ratio/Gebruik molverhouding:  $n(\text{Zn}(\text{OH})_2) : n(\text{HBr}) = 1 : 2$  ✓
- Substitution of/Vervanging van:  $99 \text{ g}\cdot\text{mol}^{-1}$  ✓  $(65 + 32 + 2)$
- Final answer/Finale antwoord:  $1,5964 \text{ g}$  (range/gebied:  $1,58 - 1,68$ ) ✓

*numbers correct.*

**POSITIVE MARKING FROM Q7.3.1/POSITIEWE NASIEN VANAF V7.3.1**

**OPTION 1/OPSIE 1**

$$n(\text{HBr})_{\text{initial/begin}} = cV$$

$$= (0,45)(0,09) \checkmark$$

$$= 0,0405 \text{ mol}$$

*0,45 and 0,09 together.*

$$n(\text{HBr reacted with/reageer met Zn}(\text{OH})_2) = 0,0405 - 0,00825 \checkmark \checkmark$$

$$= 0,03224 \text{ mol}$$

*from above.*

$$n(\text{Zn}(\text{OH})_2) = \frac{1}{2}n(\text{HBr}) = \frac{1}{2}(0,03224) \checkmark = 0,016125 \text{ mol}$$

$$m(\text{Zn}(\text{OH})_2) = nM$$

$$= (0,016125)(99) \checkmark$$

$$= 1,596 \text{ g} \checkmark$$

*65 + 2(16 + 1)*

**OPTION 2/OPSIE 2**

$$c(\text{HBr}) = 0,45 - 0,092 \checkmark \checkmark$$

$$= 0,358 \text{ mol}\cdot\text{dm}^{-3}$$

$$n(\text{HBr reacted/reageer}) = cV$$

$$= 0,358 \times 0,09 \checkmark$$

$$= 0,0322 \text{ mol}$$


$$n(\text{Zn}(\text{OH})_2) = \frac{1}{2}n(\text{HBr}) = \frac{1}{2}(0,0322) \checkmark = 0,01611 \text{ mol}$$

$$m(\text{Zn}(\text{OH})_2) = nM$$

$$= 0,01611 \times 99 \checkmark$$

$$= 1,595 \text{ g} \checkmark \quad (1,60 \text{ g})$$

(6)  
[17]



*Handwritten signatures and initials.*



**QUESTION 8/VRAAG 8**

8.1 Chemical to electrical/*Chemies na elektries* ✓ (1)

8.2 Provides path for movement of ions./ Completes the circuit./Ensures electrical neutrality in the cell./Restore charge balance. ✓  
*Verskaf pad vir beweging van ione./Voltooi die stroombaan./Verseker elektriese neutraliteit in die sel./Herstel balans van lading.* (1)

8.3 **OPTION 1/OPTION 1**

$$E_{\text{cell}}^{\ominus} = E_{\text{cathode}}^{\ominus} - E_{\text{anode}}^{\ominus} \checkmark$$

$$1,49 = 1,36 - E_{\text{anode}}^{\ominus} \checkmark$$

$$E_{\text{anode}}^{\ominus} = 1,36 - 1,49 \\ = -0,13 \text{ (V)} \checkmark$$

X is Pb/Lead/Lood ✓

**Notes/Aantekeninge**

- Accept any other correct formula from the data sheet. /Aanvaar enige ander korrekte formule vanaf gegewensblad.
- Any other formula using unconventional abbreviations, e.g.  $E_{\text{cell}}^{\ominus} = E_{\text{OA}}^{\ominus} - E_{\text{RA}}^{\ominus}$  followed by correct substitutions: /Enige ander formule wat onkonvensionele afkortings gebruik, bv.  $E_{\text{sel}}^{\ominus} = E_{\text{OM}}^{\ominus} - E_{\text{RM}}^{\ominus}$  gevolg deur korrekte vervangings:  $\frac{4}{5}$

**OPTION 2/OPSIE 2**



X is Pb/Lead/Lood ✓ *independeent*



**POSITIVE MARKING FROM Q8.3/POSITIEWE NASIEN VANAF V8.3**

8.4 X/Pb/Lead/Lood ✓  $\text{Pb} \rightarrow \text{Pb}^{2+} + 2e^-$  (1)


8.5

8.5.1 Reaction reached equilibrium./ (In each half cell) the rate of oxidation is equal to rate of reduction./Rate of the forward reaction is equal to the rate of the reverse reaction. ✓  
*Reaksie bereik ewewig./ (In elke halfsel) die tempo van oksidasie is gelyk aan tempo van reduksie./Tempo van die voorwaartse reaksie is gelyk aan die tempo van die terugwaartse reaksie.* (1)

8.5.2 Increases/Toeneem ✓ (1)

8.5.3 • [Cl<sup>-</sup>] decreases/neem af. ✓  
• Forward reaction is favoured./Voorwaartse reaksie word bevoordeel. ✓ (2)

*independeent van 1<sup>st</sup> bukket.*



*Handwritten signatures and initials*

**QUESTION 9/VRAAG 9**

9.1

**Marking guidelines/Nasienriglyne**

If any one of the underlined key phrases in the **correct context** is omitted, deduct 1 mark. *Indien enige van die onderstreepte frases in die korrekte konteks uitgelaat is, trek 1 punt af.*

The chemical process in which electrical energy is converted to chemical energy. ✓✓

*Die chemiese proses waarin elektriese energie omgeskakel word na chemiese energie.*

**OR/OF**

The use of electrical energy to produce a chemical change.

*Die gebruik van elektriese energie om 'n chemiese verandering teweeg te bring.*

**OR/OF**

The process during which an electrical current passes through a solution/molten ionic compound.

*Die proses waar 'n elektriese stroom deur 'n oplossing/gesmelte ioniese verbinding gestuur word.*


(2)

9.2




Ignore phases/Ignoreer fases

**Marking guidelines/Nasienriglyne**



• Ignore if charge omitted on electron. *Ignoreer indien lading weggelaat op elektron.*

• If charge (-) omitted on  $\text{OH}^-$  *Indien lading (-) weggelaat op  $\text{OH}^-$ :*



Max./Maks:  $\frac{1}{2}$

(2)

9.2.2 Water/  $\text{H}_2\text{O}$  ✓

(1)

9.3  $\text{H}_2\text{O}$  is a stronger oxidising agent ✓ than  $\text{Na}^+$  ✓ and will be reduced ✓ (to  $\text{H}_2$ ).  
 *$\text{H}_2\text{O}$  is 'n sterker oksideermiddel as  $\text{Na}^+$  en sal gereduseer word (na  $\text{H}_2$ ).*

**OR/OF**

$\text{Na}^+$  is a weaker oxidizing agent ✓ than  $\text{H}_2\text{O}$  ✓ and therefore  $\text{H}_2\text{O}$  will be reduced ✓ (to  $\text{H}_2$ )

*$\text{Na}^+$  is 'n swakker oksideermiddel as  $\text{H}_2\text{O}$  en daarom sal  $\text{H}_2\text{O}$  gereduseer word (na  $\text{H}_2$ )*

**OR/OF**

The half-reaction that produces  $\text{H}_2(\text{g})$  has a more positive reduction potential (-0,83 V) ✓ than the half-reaction that produces  $\text{Na}$  (-2,71 V). ✓

Therefore water/ $\text{H}_2\text{O}$  will be reduced ✓ to  $\text{H}_2$ ./ $\text{Na}^+$  will not be reduced to  $\text{Na}$ .

*Die halfreaksie wat  $\text{H}_2(\text{g})$  vorm, het 'n meer positiewe reduksiepotensiaal*

*(-0,83 V) as die halfreaksie wat  $\text{Na}$  vorm (-2,71 V).*

*Daarom word water/ $\text{H}_2\text{O}$  na  $\text{H}_2$  gereduseer./ $\text{Na}^+$  sal nie gereduseer word na  $\text{Na}$  nie.*

(3)

[8]

DEPARTMENT OF BASIC EDUCATION  
 PRIVATE BAG 8893, PRETORIA 0001  
 2019-11-13  
 APPROVED MARKING GUIDELINE  
 PUBLIC EXAMINATION

**QUESTION 10/VRAAG 10**

10.1

10.1.1 Hydrogen/Waterstof/H<sub>2</sub> ✓

10.1.2 Nitrogen monoxide/Stikstofmonoksied/NO ✓

10.1.3 Nitric acid/Salpetersuur/HNO<sub>3</sub> ✓ Hydrogen nitrate. ← is liq. already.

(4)

(1)

(1)

10.2

10.2.1 (Catalytic) oxidation/Redox/(Katalitiese) oksidasie/Redoks ✓

(1)

10.2.2 NH<sub>3</sub> + HNO<sub>3</sub> ✓ → NH<sub>4</sub>NO<sub>3</sub> ✓ Bal ✓

**Notes/Aantekeninge**


- Reactants ✓ Products ✓ Balancing ✓  
Reaktanse Produkte Balansering
- Ignore double arrows (⇌) and phases./Ignoreer dubbelpyle (⇌) en fases.
- Marking rule 6.3.10./Nasiennreël 6.3.10. No arrow, no marks. ✓

(3)

10.3

10.3.1 (Total) percentage of nutrients/fertiliser/N,P,K. ✓  
(Totale) persentasie nutriente/ kunsmis/N,P, K.

(1)



Handwritten signatures and initials at the bottom right of the page.

10.3.2 **Marking guidelines/Nasiernriglyne**

- Calculate mass fertiliser in A./Bereken massa kunsmis in A ✓
- Calculate mass fertiliser in B./ Bereken massa kunsmis in B ✓
- Calculate mass P in A and B ./Bereken massa P in A en B ✓
- Final answer/Finale antwoord: **Independent from above.**  
 B has more phosphorous than/het meer fosfor as A. ✓

<p><b>OPTION 1/OPSIE 1</b>                  Mass fertiliser in A:                  Massa kunsmis in A:  <math>m = \frac{21}{100} \times 50 \checkmark = 10,5 \text{ kg}</math>                  Mass fertiliser in B:                  /Massa kunsmis in B:  <math>m = \frac{27}{100} \times 40 \checkmark = 10,8 \text{ kg}</math>                  Mass phosphorous in A/                  Massa fosfor in A:  <math>\frac{3}{8} \times 10,5 = 3,94 \text{ kg}</math>                  Mass phosphorous in B/                  Massa fosfor in B:  <math>\frac{3}{8} \times 10,8 = 4,05 \text{ kg}</math>                  Fertiliser B has more phosphorous than fertiliser A. ✓</p>	<p><b>OPTION 3/OPSIE 3</b>                  Mass phosphorous in A/                  Massa fosfor in A:  <math>\%P = \frac{3}{8} \times 21 = 7,88\%</math>  <math>m(P) = \frac{7,88}{100} \times 50 \checkmark = 3,94 \text{ kg}</math>                  Mass(P) in B                  Massa (P) in B:  <math>\%(P) = \frac{3}{8} \times 27 = 10,13\%</math>  <math>m = \frac{10,13}{100} \times 40 \checkmark = 4,05 \text{ kg}</math>                  Fertiliser B has more phosphorous than fertiliser A. /Kunsmis B het meer fosfor as kunsmis A ✓</p>
<p><b>OPTION 2/OPSIE 2</b>                  Mass phosphorous in A/                  Massa fosfor in A:  <math>m = \frac{3}{8} \times \frac{21}{100} \times 50 \checkmark = 3,94 \text{ kg}</math>                  Mass(P) in B                  Massa (P) in B:  <math>m = \frac{3}{8} \times \frac{27}{100} \times 40 \checkmark = 4,05 \text{ kg}</math>                  Fertiliser B has more phosphorous than fertiliser A. /Kunsmis B het meer fosfor as kunsmis A. ✓</p>	<p><b>OPTION 4/OPSIE 4</b>                  Mass fertiliser in A:                  Massa kunsmis in A:  <math>m = \frac{21}{100} \times 50 \checkmark = 10,5 \text{ kg}</math>                  Mass fertiliser in B:                  /Massa kunsmis in B:  <math>m = \frac{27}{100} \times 40 \checkmark = 10,8 \text{ kg}</math>                  For the same NPK ratio ✓                  the bag with more fertiliser will have more phosphorous ∴ bag B ✓                  Vir dieselfde NPK verhouding, die sake met meer kunsmis sal meer fosfor het ∴ sak B</p>

OR A has less than B.

**DEPARTMENT OF BASIC EDUCATION**  
 PRIVATE BAG X899, PRETORIA 0001  
 2019 -11- 13  
**APPROVED MARKING GUIDELINE**  
 PUBLIC EXAMINATION

(4)  
 [12]  
**TOTAL/TOTAAL: 150**

*Handwritten signatures and initials*